

n -dimensional local fields and adeles on n -dimensional schemes.

D.V. Osipov *

1 Introduction

The notion of n -dimensional local field has appeared in the works of A. N. Parshin and K. Kato in the middle of 70's. These fields generalize the usual local fields (which is 1-dimensional in this sense) and help us to see on higherdimensional algebraic schemes from the local point of view.

With every flag

$$X_0 \subset X_1 \dots \subset X_{n-1}$$

$$\dim X_i = i$$

of irreducible subvarieties on a scheme X ($\dim X = n$) one can canonically associate a ring $K_{(X_0, \dots, X_{n-1})}$. In case everything is regularly embedded, the ring is an n -dimensional local field.

Originally, higherdimensional local fields were used for the development of generalisation of class field theory to the schemes of arbitrary dimension (works of A. N. Parshin, K. Kato, S. V. Vostokov and others), [22], [11]. But many problems of algebraic varieties can be reformulated in terms of higherdimensional local fields and higher adelic theory.

For a scheme X adelic object

$$\mathbb{A}_X = \prod' K_{(X_0, \dots, X_{n-1})}$$

where the product is taken over all the flags with respect to certain restrictions on components of adeles. A.N. Parshin defined in [19] adeles on algebraic surfaces, which generalize usual adeles on curves. A.A. Beilinson introduced a simplicial approach to adeles and generalized to arbitrary dimensional Noetherian schemes in [2].

A. N. Parshin, A. A. Beilinson, A. Huber, A. Yekutieli, V. G. Lomadze and others described connections of higher adelic groups with cohomology of coherent sheaves ([19], [2], [8], [29], [3], [4]), intersection theory ([21], [12], [15], [4]), Chern classes ([21], [10], [4]), theory of residues ([19], [29], [2], [13], [4]), torus actions ([5]).

*The author was supported by an LMS grant for young Russian mathematicians at the University of Manchester, and also by the Russian Foundation for Basic Research, grant no. 05-01-00455.

This paper is a survey of basic notions of higherdimensional local fields and adeles on higherdimensional schemes.

The paper is organized as follows.

In section 2 we give a general definition of n -dimensional local field and formulate classification theorems of n -dimensional local fields. We describe how n -dimensional local fields appear from algebraic varieties and arithmetical schemes.

In section 3 we define higher dimensional adeles and adelic complexes. Starting from an example of adelic complexes on algebraic curves, we give a general simplicial definition for arbitrary Noetherian schemes, which is due to A.A. Beilinson. We formulate the theorems about adelic resolutions of quasicoherent sheaves on Noetherian schemes. We applicate this general constructions to algebraic surfaces to obtain adelic complexes on algebraic surfaces, which were introduced by A.N. Parshin.

In section 4 we describe restricted adelic complexes. In constrast to the adelic complexes from section 3, restricted adelic complexes are connected with a single flag of subvarieties. A.N. Parshin introduced restricted adeles for algebraic surfaces in [24], [23]. The author introduced restricted adelic complexes for arbitrary schemes in [17]. We give also the reconstruction theorem on restricted adelic complexes.

In the last section we briefly describe reciprocity laws on algebraic surfaces.

The author is very grateful to A.N. Parshin for a lot of discussions on higherdimensional local fields and adeles. The author is also grateful to M. Taylor for interesting discussions and his hospitality during the visit to Manchester sponsored by an LMS grant.

2 n -dimensional local fields

2.1 Classification theorems.

We fix a perfect field k .

We say that K is a local field of dimension 1 with the residue field k , if K is a fraction field of complete discrete valuation ring $\mathcal{O}_K$ with the residue field $\bar{K} = k$.

We denote by ν_K the discrete valuation of K and by m_K the maximal ideal of ring $\mathcal{O}_K$.

Such a field has the following structure

$$K \supset \mathcal{O}_K \rightarrow \bar{K} = k$$

.

As examples of such fields we have the field of power series

$$K = k((t)), \quad \mathcal{O}_K = k[[t]], \quad \bar{K} = k$$

and the field of p-adic numbers

$$K = \mathbb{Q}_p, \quad \mathcal{O}_K = \mathbb{Z}_p, \quad k = \mathbb{F}_p.$$

Moreover, we have only the following possibilities, see [27, ch. II]:

Theorem 1 *Let K be a local field of dimension 1 with the residue field k then*

1. $K = k((t))$ is the power series field if $\text{char} K = \text{char} k$;
2. (a) $K = \text{Frac}(W(k))$ where $\mathcal{O}_K = W(k)$ is the Witt ring (for example, $K = \mathbb{Q}_p$),
(b) or K is a finite totally ramified extension of the field $\text{Frac}(W(k))$
if $\text{char} K = 0$, $\text{char} k = p$.

Now we give the following inductive definition.

Definition 1 *We say that a field K is a local field of dimension n with the last residue field k if*

1. $n = 0$ and $K = k$
2. $n \geq 1$ and K is the fraction field of a complete discrete valuation ring $\mathcal{O}_K$ whose residue field $\bar{K}$ is a local field of dimension $n - 1$ with the last residue field k

A local field of dimension $n \geq 1$ has the following inductive structure:

$$K = K^{(0)} \supset \mathcal{O}_K \rightarrow \bar{K} = K^{(1)} \supset \mathcal{O}_{\bar{K}} \rightarrow \bar{\bar{K}}^{(1)} = K^{(2)} \supset \mathcal{O}_{K^{(2)}} \rightarrow \dots \bar{K}^{(n)} = k,$$

where for a discrete valuation field F $\mathcal{O}_F$ the ring of integers in F and $\bar{F}$ the residue field. The maximal ideals in $\mathcal{O}_{K^{(i)}}$ we denote by $m_{K^{(i)}}$.

Every field $K^{(i)}$ is a local field of dimension $n - i$ with the last residue field k .

Definition 2 *A collection of elements $t_1, \dots, t_n \in \mathcal{O}_K$ is called a system of local parameters, if for all $i = 1, \dots, n$*

$$t_i \bmod m_{K^{(0)}} \in \mathcal{O}_{K^{(1)}}, \dots, t_i \bmod m_{K^{(i-2)}} \in \mathcal{O}_{K^{(i-1)}}$$

and the element $t_i \bmod m_{K^{(i-2)}} \in \mathcal{O}_{K^{(i-1)}}$ is a generator of $m_{K^{(i-1)}}$.

An example of an n -dimensional local field is the field $K = k((t_n)) \dots ((t_1))$. For this field we have

$$K^{(i)} = k((t_n)) \dots ((t_{i+1})) \quad , \quad \mathcal{O}_{K^{(i)}} = k((t_n)) \dots ((t_{i+2}))[t_{i+1}].$$

We consider $n = 2$, then we have

$$K \supset \mathcal{O}_K \rightarrow \bar{K} \supset \mathcal{O}_{\bar{K}} \rightarrow k.$$

We can construct the following examples of 2-dimensional local fields with the last residue field k . These examples depend on characteristic of 2-dimensional field K and characteristics of its residue fields.

- $K = k((t_2))((t_1))$. And $\text{char} K = \text{char} \bar{K} = \text{char} k$.

- $K = F((t))$, where F is a local field of dimension 1 with the residue field k such that $\text{char}(F) \neq \text{char}(k)$, for example $F = \mathbb{Q}_p$.
- $K = F\{\{t\}\}$, where F is a local field of dimension 1 with the residue field k .

The field $F\{\{t\}\}$ has the following description

$$a \in F\{\{t\}\} \quad : \quad a = \sum_{i=-\infty}^{i=+\infty} a_i t^i, \quad a_i \in F,$$

where $\lim_{i \rightarrow -\infty} \nu_F(a_i) = 0$ and $\nu_F(a_i) > c_a$ for some integer c_a .

We put the discrete valuation $\nu_{F\{\{t\}\}}(a) = \min \nu_F(a_i)$. Then the ring $\mathcal{O}_{F\{\{t\}\}}$ consists of elements a such that all $a_i \in \mathcal{O}_F$, and the maximal ideal $m_{F\{\{t\}\}}$ consists of elements a such that all $a_i \in m_F$. Therefore

$$\overline{F\{\{t\}\}} = \bar{F}((t)).$$

We remark that for $F = k((u))$ the field $F\{\{t\}\}$ is isomorphic to the field $k((t))((u))$. There is the following classification theorem, see [4], [20], [30], [31].

Theorem 2 *Let K be an n -dimensional local field with the finite last residue field. Then*

1. K is isomorphic to $\mathbb{F}_q((t_n)) \dots ((t_1))$ if $\text{char}(K) = p$;
2. K is isomorphic to $F((t_{n-1})) \dots ((t_1))$, F is a 1-dimensional local field, if $\text{char}(K^{(n-1)}) = 0$.
3. K is a finite extension of a field

$$F\{\{t_n\}\} \dots \{\{t_{m+2}\}\}((t_m)) \dots ((t_1)) \quad (*)$$

and there is a finite extension of K which is of the form $(*)$, but possibly with different F and t_i , if $\text{char}(K^{(m)}) = 0$, $\text{char}(K^{(m+1)}) = p$.

We remark that if π is a local parameter for a 1-dimensional local field F then $t_1, \dots, t_m, \pi, t_{m+2}, \dots, t_n$ are local parameters for a field

$$F\{\{t_n\}\} \dots \{\{t_{m+2}\}\}((t_m)) \dots ((t_1)).$$

2.2 Local fields which come from algebraic geometry.

We consider an algebraic curve C over the field k . We fix a smooth point $p \in C$. We consider

$$K_p = \text{Frac}(\hat{\mathcal{O}}_p),$$

where

$$\hat{\mathcal{O}}_p = \varprojlim_n \mathcal{O}_p / m_p^n$$

is a completion of the local ring $\mathcal{O}_p$ of the point p on the curve C . Then

$$K_p = k(p)((t)), \quad (1)$$

where $k(p) = \mathcal{O}_p/m_p$ is the residue field of the point p on the curve C , which is the finite extension of the field k , and t is a local parameter of the point p on the curve C .

We see that the field K_p corresponds to the case 1 of the classification theorem 1.

Now we consider a field of algebraic numbers K , which is a finite extension of the field $\mathbb{Q}_p$. We consider the ring of integers A of the field K . Let $X = \text{Spec } A$ be a 1-dimensional scheme. We fix a closed point $p \in X$, which corresponds to a maximal ideal in A . Then the completion of the field K at the point p is

$$K_p = \text{Frac} \left(\varprojlim_n A_p/m_p^n \right) \quad (2)$$

We see that K_p is a 1-dimensional local field with the residue field $\mathbb{F}_q$, and the field K_p corresponds to the case 2 of the classification theorem 1.

We give now the definitions for the general situation.

Let R be a ring, p a prime ideal of R , M an R -module.

Let $S_p = R \setminus p$. We write $S_p^{-1}M$ for the localisation of M at S_p .

For an ideal a of R set $C_a M = \varprojlim_{n \in \mathbb{N}} M/a^n M$.

Let X be a Noetherian scheme of dimension n . Let $\delta = (p_0, \dots, p_n)$ be a chain of points of X (i.e. the chain of integral irreducible subschemes if we consider the closures of points p_i) such that $p_{i+1} \in \overline{\{p_i\}}$ for any i ($\overline{\{p_i\}}$ is the closure of the point p_i in X). We suppose that for all i $\dim p_i = i$. We restrict δ to some affine open neighbourhood $\text{Spec } B$ of the closed point p_n on X . Then δ determines a chain of prime divisors of the ring B , which we denote by the same letters $(p_0, \dots, p_n)$. We define a ring

Definition 3

$$K_\delta \stackrel{\text{def}}{=} C_{p_0} S_{p_0}^{-1} \dots C_{p_n} S_{p_n}^{-1} B \quad (3)$$

This definition of K_δ does not depend on the choice of affine neighbourhood $\text{Spec } B$ of the point p_n on the scheme X , see [8, prop.3.1.3., prop. 3.2.1.].

We remark that the ring $C_{p_n} S_{p_n}^{-1} B$ from formula 3 coincides with the completion $\hat{\mathcal{O}}_{p_n, X}$ of the local ring of the point p_n on the scheme X .

We consider now examples of formula (3) for small n .

Example 1 If X is an irreducible 1-dimensional scheme (X is an irreducible curve over the field k or the spectrum of the ring of algebraic integers), p is a smooth point of X , η is a general point of X , then for $\delta = (\eta, p)$ we obtain that K_δ is a 1-dimensional local field, which coincides with the field K_p from formula (1) or (2).

Example 2 Now let X be an irreducible algebraic surface over the field k . Let C be an irreducible divisor on X and p a point on C . We suppose that p is a smooth point on X and on C . Let η be a general point of X . We consider $\delta = (\eta, C, p)$. We fix the local parameter $t \in k(X)$ of the curve C on X at the point p ($t = 0$ is a local equation of the curve C at the point p on X), and local parameter $u \in k(X)$ at the point p on X which is transversal to the local parameter t (the divisor $u = 0$ is transversal to the divisor $t = 0$ at the point p). We fix any affine neighbourhood $\text{Spec } B$ of p on X . Then

$$\begin{aligned} C_p S_p^{-1} B &= k(p)[[u, t]] \\ C_C S_C^{-1} C_p S_p^{-1} B &= k(p)((u))[[t]] \\ K_\delta &= C_\eta S_\eta^{-1} C_C S_C^{-1} C_p S_p^{-1} B = k(p)((u))((t)). \end{aligned}$$

Hence K_δ is a 2-dimensional local field with the last residue field $k(p)$.

We see that the field K_δ corresponds to the case 1 of the classification theorem 2.

Example 3 The previous example can be generalized. Let $p_0, \dots, p_n$ be a flag of irreducible subvarieties on n -dimensional algebraic variety X over the field k such that $\dim p_i = n - i$, $p_{i+1} \subset p_i$ for all i and the point p_n is a smooth point on all subvarieties p_i . We can choose a system of local parameters $t_1, \dots, t_n \in \mathcal{O}_{p_n, X}$ of the point p_n on X such that for every i equations $t_1 = 0, \dots, t_i = 0$ define a subvariety p_i in some neighbourhood of the point p_n on X . Then according to formula (3) and similar to the previous example we have for $\delta = (p_0, \dots, p_n)$

$$K_\delta = k(p)((t_n)) \dots ((t_1)).$$

Example 4 Now we suppose that a scheme X is an arithmetical surface, i.e., $\dim X = 2$ and we have a flat, projective morphism $f : X \rightarrow Y = \text{Spec } A$, where A is the ring of integers of a number field K . We consider two kinds of integral irreducible 1-dimensional closed subschemes C on X .

1. A subscheme C is horizontal, i.e., $f(C) = Y$. We consider a point $x \in C$ which is smooth on X and C . Let $\delta = (\eta, C, x)$, where η is a general point of X . Then

$$K_\delta = L((t)),$$

where $t = 0$ is a local equation of C at the point x on X and $L \supset K_{f(x)} \supset \mathbb{Q}_p$ is a finite extension. Thus, K_δ is a 2-dimensional local field with the finite last residue field.

We see that this field K_δ corresponds to the case 2 of the classification theorem 2.

2. A subscheme C is vertical, i.e. it is a component of a fibre of f . This C is defined over some finite field $\mathbb{F}_q$. We consider a point $x \in C$ such that the morphism f is smooth at x and the point x is also defined over the field $\mathbb{F}_q$. Let $\delta = (\eta, C, x)$, where η is a general point of X . Then we apply formula (3). For any affine neighbourhood $\text{Spec } B$ of x on X the ring $C_x S_x^{-1} B$ coincides with the completion

$\hat{\mathcal{O}}_{x,X}$ of the local ring of the point x on X . But since f is a smooth map at p , the ring

$$\hat{\mathcal{O}}_{x,X} = \mathcal{O}_{K_{f(x)}}[[u]].$$

Therefore we obtain

$$K_\delta = K_{f(x)}\{\{u\}\},$$

where $K_{f(x)} \supset \mathbb{Q}_p$ is a finite extension. Thus, K_δ is a 2-dimensional local field with the finite last residue field.

We see that this field K_δ corresponds to the case 3 of the classification theorem 2.

We remark that in these both cases we have a canonical embedding f^* of the 1-dimensional local field $K_{f(x)}$ to the 2-dimensional local field K_δ .

Now we consider only excellent Noetherian schemes X (e.g. a scheme of finite type over a field, over $\mathbb{Z}$, or over a complete semi-local Noetherian ring; see [14, §34] and [6, §7.8]).

We introduce the following notations (see [21]). Let $\delta = (p_0, \dots, p_n)$. Let a subscheme $X_i = \overline{\{p_i\}}$ be the closure of the point p_i in X . We introduce by induction the schemes X'_{i,α_i} in the following diagramm

$$\begin{array}{ccccccc} X_0 & \supset & X_1 & \supset & X_2 & \supset & \dots \\ \uparrow & & \uparrow & & & & \\ X'_0 & \supset & X_{1,\alpha_1} & \supset & \uparrow & & \\ & & \uparrow & & & & \\ & & X'_{1,\alpha_1} & \supset & X_{2,\alpha_2} & & \\ & & & & \uparrow & & \\ & & & & \vdots & & \end{array}$$

Here X' is the normalization of a scheme x and X_{i,α_i} is an integral irreducible subscheme in $X'_{i-1,\alpha_{i-1}}$ which is mapped onto X_i . Then by any such diagram we obtain the collection of indices $(\alpha_1, \dots, \alpha_n)$. We denote the finite set of all such collections of indices by Λ_δ .

Such a collection of indices $(\alpha_1, \dots, \alpha_n) \in \Lambda_\delta$ determines a chain of discrete valuations in the following way. Integral irreducible subvariety X_{1,α_1} of the normal scheme X'_0 defines the discrete valuation of the field of functions on X_0 . The residue field of this discrete valuation is the field of functions on the normal scheme X'_{1,α_1} , and the integer irreducible subscheme X_{2,α_2} defines the discrete valuation here. We have to proceed further for $\dots, \alpha_3, \dots, \alpha_n$ in this way.

Moreover, there is the following theorem ([21]). (See also [29, theorem 3.3.2] for the proof).

Theorem 3 *Let X be an integral excellent n -dimensional Noetherian scheme. Then for $\delta = (p_0, \dots, p_n)$ the ring K_δ is an Artinian ring and*

$$K_\delta = \prod_{(\alpha_1, \dots, \alpha_n) \in \Lambda_\delta} K_{(\alpha_1, \dots, \alpha_n)} \quad ,$$

where every $K_{(\alpha_1, \dots, \alpha_n)}$ is an n -dimensional local field.

Example 5 To illustrate this theorem we compute now the ring K_δ in the following situation. Let p be a smooth point on an irreducible algebraic surface X over k . Suppose an irreducible curve $C \subset X$ contains the point p , but C has at the point p the node singularity, i.e., completed local ring of the point p on the curve C is $k[[t, u]]/tu$ for some local formal parameters u, t of the point p on X . Let $\delta = (\eta, C, p)$, where η is a general point of X . We fix any affine neighbourhood $\text{Spec } B$ of p on X . Then according to formula (3)

$$\begin{aligned} C_p S_p^{-1} B &= k(p)[[u, t]] \\ C_C S_C^{-1} C_p S_p^{-1} B &= k(p)((u))[[t]] \oplus k(p)((t))[[u]] \\ K_\delta = C_\eta S_\eta^{-1} C_C S_C^{-1} C_p S_p^{-1} B &= k(p)((u))((t)) \oplus k(p)((t))((u)). \end{aligned}$$

3 Adeles and adelic complexes

3.1 Adeles on curves

Let C be a smooth connected algebraic curve over the field k .

For any coherent sheaf $\mathcal{F}$ on C we consider an adelic space $\mathbb{A}_C(\mathcal{F})$:

$$\mathbb{A}_C(\mathcal{F}) = \left\{ \{f_p\} \in \prod_{p \in C} \mathcal{F} \otimes_{\mathcal{O}_C} K_p \text{ such that } f_p \in \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p} \text{ for almost all } p, \right\}$$

where the product is over all closed points p of the curve C .

We construct the following complex $\mathcal{A}_C(\mathcal{F})$:

$$\begin{array}{ccccc} \mathcal{F} \otimes_{\mathcal{O}_C} k(C) & \times & \prod_{p \in C} \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p} & \longrightarrow & \mathbb{A}_C(\mathcal{F}) \\ a & \times & b & \mapsto & a + b. \end{array}$$

We have the following theorem (for example, see [26]).

Theorem 4 *The cohomology groups of the complex $\mathcal{A}_C(\mathcal{F})$ coincide with the cohomology $H^*(C, \mathcal{F})$, where $\mathcal{F}$ is any coherent sheaf on C .*

Proof. We give here the sketch of the proof. We write an adelic complex $\mathcal{A}_U(\mathcal{F})$ of the sheaf $\mathcal{F}$ for any open subset $U \subset C$. Taking into account all U we obtain a complex of sheaves $\mathcal{A}(\mathcal{F})$ on the curve C . Then for the small affine U we obtain that the following complex

$$0 \longrightarrow \mathcal{F}(U) \longrightarrow \mathcal{A}_U(\mathcal{F}) \longrightarrow 0$$

is exact, since we can apply the approximation theorem for Dedekind rings over fields. Therefore the complex $\mathcal{A}(\mathcal{F})$ is a resolution of the sheaf $\mathcal{F}$ on C . And by construction this resolution is a flasque resolution of the sheaf $\mathcal{F}$ on C . Therefore it calculates the cohomology of the sheaf $\mathcal{F}$ on the curve C .

3.2 Adeles on higherdimensional schemes

In this section we give a generalization of adelic complexes to the schemes of arbitrary dimensions.

For algebraic surfaces adelic complexes were introduced by A.N. Parshin in [19]. We will give a detail exposition of adelic complexes on algebraic surfaces later as an application of general machinery, which was constructed for arbitrary Noetherian schemes by A.Beilinson in [2]. For a good exposition and proofs of Beilinson results see [8].

3.2.1 Definition of adelic spaces.

We introduce the following notations. For any Noetherian scheme X let $P(X)$ be the set of points of the scheme X . Consider $p, q \in P(X)$. Define $p \geq q$ if $q \in \overline{\{p\}}$, i.e., the point p is in closure of the point q . Then $\geq$ is a half ordering on $P(X)$. Let $S(X)$ be the simplicial set induced by $(P(X), \geq)$, i.e.,

$$S(X)_m = \{(p_0, \dots, p_m) \mid p_i \in P(X); p_i \geq p_{i+1}\}$$

is the set of m -simplices of $S(X)$ with the usual boundary δ_i^n and degeneracy maps σ_i^n for $n \in \mathbb{N}$, $0 \leq i \leq n$.

Let $K \subset S(X)_n$. For $p \in P(V)$ we denote

$$_p K = \{(p_1 > \dots > p_n) \in S(V)_{n-1} \mid (p > p_1 > \dots > p_n) \in K\}.$$

Let $\mathbf{QS}(X)$ and $\mathbf{CS}(X)$ be the category of quasicoherent and coherent sheaves on the scheme X . Let $\mathbf{Ab}$ be the category of Abelian groups.

We have the following proposition, see [2], [8], [9], which is also a definition.

Proposition 1 *Let $S(X)$ be the simplicial set associated to the Noetherian scheme X . Then there exist for integer $n > 0$, $K \in S(X)_n$ functors*

$$\mathbb{A}(K, \cdot) : \mathbf{QS}(X) \longrightarrow \mathbf{Ab}$$

uniquely determined by the properties 1, 2, 3, which are additive and exact.

1. $\mathbb{A}(K, \cdot)$ commutes with direct limits.
2. For $n = 0$, a coherent sheaf $\mathcal{F}$ on X

$$\mathbb{A}(K, \mathcal{F}) = \prod_{p \in K} \varprojlim_l \mathcal{F}_p / m_p^l \mathcal{F}_p.$$

3. For $n > 0$, a coherent sheaf $\mathcal{F}$ on X

$$\mathbb{A}(K, \mathcal{F}) = \prod_{p \in P(V)} \varprojlim_l \mathbb{A}({}_\eta K, \mathcal{F}_p / m_p^l \mathcal{F}_p)$$

Remark 1 Since any quasicoherent sheaf on an excellent Noetherian scheme is a direct limit of coherent sheaves, we can apply property 1 of this proposition to define $\mathbb{A}(K, \mathcal{F})$ on quasicoherent sheaves.

3.2.2 Local factors.

The definition of $\mathbb{A}(K, \mathcal{F})$ is a definition of inductive kind. By induction of the definition there is the following proposition, [8, prop. 2.1.4.].

Proposition 2 *For integer $n > 0$, $K \subset S(X)_n$, a quasicoherent sheaf $\mathcal{F}$ on X*

$$\mathbb{A}(K, \mathcal{F}) \subset \prod_{\delta \in K} \mathbb{A}(\delta, \mathcal{F}).$$

The inclusion is a transformation of functors.

From this proposition we see that $\mathbb{A}(K, \mathcal{F})$ is a kind of complicated adelic product inside of $\prod_{\delta \in K} \mathbb{A}(\delta, \mathcal{F})$.

Therefore it is important to study the local factors $\mathbb{A}(\delta, \mathcal{F})$ for $\delta \in S(X)_n$. We have the following two propositions from [8] about these local factors..

Proposition 3 *Let $\delta = (p_0, \dots, p_n) \in S(X)_n$. Let U be an open affine subscheme which contains the point p_n and therefore all of δ . Let $M = \mathcal{F}(U)$. Then for a quasicoherent sheaf $\mathcal{F}$*

$$\mathbb{A}(\delta, \mathcal{F}) = \mathbb{A}(\delta, \tilde{M}),$$

where $\tilde{M}$ is a quasicoherent sheaf on affine U which corresponds to M .

In the following proposition local factors $\mathbb{A}(\delta, \mathcal{F})$ are computed for affine schemes.

Proposition 4 *Let $X = \text{Spec } R$ and $\mathcal{F} = \tilde{M}$ for some R -module M . Further let $\delta \in (p_0, \dots, p_n) \in S(X)_n$. Then*

$$\mathbb{A}(\delta, \mathcal{F}) = C_{p_0} S_{p_0}^{-1} \dots C_{p_n} S_{p_n}^{-1} R \otimes_R M. \quad (4)$$

$C_{p_0} S_{p_0}^{-1} \dots C_{p_n} S_{p_n}^{-1} R$ is a flat Noetherian R -algebra. And for finitely generated R -modules

$$C_{p_0} S_{p_0}^{-1} \dots C_{p_n} S_{p_n}^{-1} R \otimes_R M = C_{p_0} S_{p_0}^{-1} \dots C_{p_n} S_{p_n}^{-1} M.$$

We compare now formula (4) from the last proposition and formula (3) for K_δ . We obtain that for an n -dimensional Noetherian scheme X , for $\delta \in S(X)_n$ and a quasicoherent sheaf $\mathcal{F}$

$$\mathbb{A}(\delta, \mathcal{F}) = K_\delta \otimes_{\mathcal{O}_X} \mathcal{F}.$$

Remark 2 Due to theorem 3 it means that for $\delta \in S(X)_n$ local factors $\mathbb{A}(\delta, \mathcal{O}_X)$ on excellent Noetherian integral n -dimensional scheme X are finite products of n -dimensional local fields.

3.2.3 Adelic complexes.

Now we want to define adelic complexes on the scheme X .

We have the simplicial set $S(X)$ with the usual boundary maps δ_i^n and degeneracy maps σ_i^n for $n \in \mathbb{N}$, $0 \leq i \leq n$.

We remark the following property, see [8, prop. 2.1.5.].

Proposition 5 *Let $K, L, M \subset S(X)_n$ such that $K \cup M = L$, $K \cap M = \emptyset$. Then there are natural transformations i and π of functors*

$$i(\cdot) : \mathbb{A}(K, \cdot) \longrightarrow \mathbb{A}(L, \cdot)$$

$$\pi(\cdot) : \mathbb{A}(L, \cdot) \longrightarrow \mathbb{A}(M, \cdot)$$

such that the following diagram is commutative and has split-exact lines for all quasicoherent sheaves $\mathcal{F}$ on X

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{A}(K, \mathcal{F}) & \xrightarrow{i(\mathcal{F})} & \mathbb{A}(L, \mathcal{F}) & \xrightarrow{\pi(\mathcal{F})} & \mathbb{A}(M, \mathcal{F}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \prod_{\delta \in K} \mathbb{A}(\delta, \mathcal{F}) & \longrightarrow & \prod_{\delta \in L} \mathbb{A}(\delta, \mathcal{F}) & \longrightarrow & \prod_{\delta \in M} \mathbb{A}(\delta, \mathcal{F}) \longrightarrow 0. \end{array}$$

This proposition is proved by induction on definition-proposition 1.

Definition 4 *Let $K \subset S(X)_0$, $\mathcal{F}$ a quasicoherent sheaf. Then let*

$$d^0(K, \mathcal{F}) : \Gamma(X, \mathcal{F}) \longrightarrow \mathbb{A}(K, \mathcal{F})$$

be the canoical map, which is a natural transformation of functors.

Definition 5 *Let $K \subset S(X)_{n+1}$, $L \subset S(X)_n$, $\delta_i^{n+1}K \subset L$ for some $i \in \{0, \dots, n+1\}$. We define transformations of functors*

$$d_i^{n+1}(K, L, \cdot) : \mathbb{A}(L, \cdot) \longrightarrow \mathbb{A}(K, \cdot)$$

by the following properties.

1. *If $i = 0$ and $\mathcal{F}$ is a coherent sheaf on X , then we apply the functor $\mathbb{A}({}_pK, \cdot)$ to $\mathcal{F} \rightarrow \mathcal{F}_p/m_p^l \mathcal{F}_p$ and compose this map with the projection of proposition 5 for $L \supset {}_pK$. We use the universal property of $\prod_{p \in P(X)} \lim_{\leftarrow}$.*
2. *If $i = 1$, $n = 0$ and $\mathcal{F}$ is a coherent sheaf on V , then the projection of proposition 5 for $L \supset {}_pK$ is composed with the following map. The maps $d^0({}_pK, \mathcal{F}_p/m_p^l \mathcal{F}_p)$ form a projective system for $l \in \mathbb{N}$ and we apply $\prod_{p \in P(X)} \lim_{\leftarrow}$ to it.*
3. *If $i > 0$, $n > 0$, $\mathcal{F}$ is a coherent sheaf, then the hypothesis $\delta_i^{n+1}K \subset L$ implies $\delta_{i-1}^n({}_pK) \subset {}_pL$ for all $p \in P(X)$. Set*

$$d_i^{n+1}(K, L, \mathcal{F}) = \prod_{p \in P(X)} \lim_{\leftarrow, l \in \mathbb{N}} d_{i-1}^n({}_pK, {}_pL, \mathcal{F}_p/m_p^l \mathcal{F}_p).$$

4. $d_i^{n+1}(K, L, \cdot)$ commutes with direct limits.

For $\delta \in S(V)_{n+1}$ and $\delta' = \delta_i^{n+1}(\delta) \in S(V)_n$ by definition 5 we have local boundary map

$$d_i^{n+1} : \mathbb{A}(\delta', \mathcal{F}) \longrightarrow \mathbb{A}(\delta, \mathcal{F}).$$

For $K \subset S(X)_{n+1}$, $L \subset S(X)_n$ with $\delta_i^{n+1}K \subset L$ we define

$$D_i^{n+1}(\mathcal{F}) : \prod_{\delta \in L} \mathbb{A}(\delta, \mathcal{F}) \longrightarrow \prod_{\delta \in K} \mathbb{A}(\delta, \mathcal{F}),$$

where $(x_\delta)_{\delta \in L} \mapsto (y_\delta)_{\delta \in K}$ is given by $y_\delta = d_i^{n+1}(x_{\delta'})$.

For the computations of boundary maps it is usefull the following proposition, which describes the boundary maps d_i^{n+1} by means of the boundary maps D_i^{n+1} on the product of local factors.

Proposition 6 *Let $K \subset S(X)_{n+1}$, $L \subset S(X)_n$ with $\delta_i^{n+1}K \subset L$. The following diagram commutes*

$$\begin{array}{ccc} \mathbb{A}(L, \mathcal{F}) & \xrightarrow{d_i^{n+1}} & \mathbb{A}(K, \mathcal{F}) \\ \downarrow & & \downarrow \\ \prod_{\delta \in L} \mathbb{A}(\delta, \mathcal{F}) & \xrightarrow{D_i^{n+1}} & \prod_{\delta \in K} \mathbb{A}(\delta, \mathcal{F}). \end{array}$$

This porposition is proved by induction of definitions.

Now for the scheme X we consider the set $S(X)_n^{(red)}$ of non degenerate n -dimensional simplices. (A simplex $(p_0, \dots, p_n)$ is nondegenerate if $p_i \neq p_{i+1}$ for any i .) For any $n \geq 0$, for any quasicohherent sheaf $\mathcal{F}$ on X we denote

$$\mathbb{A}_X^n(\mathcal{F}) = \mathbb{A}(S(X)_n^{(red)}, \mathcal{F}).$$

We consider the boundary maps

$$d_i^{n+1} : \mathbb{A}_X^n(\mathcal{F}) \longrightarrow \mathbb{A}_X^{n+1}(\mathcal{F}).$$

There are the following equalities for these boundary maps:

$$d_j^n d_i^n = d_i^n d_{j-1}^n \quad i < j. \quad (5)$$

For $n \geq 1$ we define $d_n : \mathbb{A}_X^{n-1}(\mathcal{F}) \longrightarrow \mathbb{A}_X^n(\mathcal{F})$ by

$$d^n = \sum_{j=0}^n (-1)^j d_j^n. \quad (6)$$

We have the following proposition, which is also a definition.

Proposition 7 *Differentials d^n make $\mathbb{A}_X^*(\mathcal{F})$ into a cohomological complex of Abelian groups $\mathcal{A}_X(\mathcal{F})$, which we call the adelic complex of the sheaf $\mathcal{F}$ on X .*

Proof. It follows by direct calculations with formulas (5) and (6).

We have the following theorem.

Theorem 5 *For any quasicoherent sheaf $\mathcal{F}$ on a Noetherian scheme X*

$$H^i(\mathcal{A}_X(\mathcal{F})) = H^i(X, \mathcal{F}).$$

Proof. The proof of this theorem is a very far generalization of the proof of theorem 4. Indeed, for any open subscheme $U \subset X$ we consider the following complex

$$0 \longrightarrow \mathcal{F}(U) \xrightarrow{d^0} \mathbb{A}_U^0(\mathcal{F}) \xrightarrow{d^1} \mathbb{A}_U^1(\mathcal{F}) \xrightarrow{d^2} \dots \xrightarrow{d^n} \mathbb{A}_U^n(\mathcal{F}) \xrightarrow{d^{n+1}} \dots \quad (7)$$

Taking into account all U , we obtain that this complex is a complex of sheaves on X . Moreover, by proposition 5 the sheaves in this complex are flasque sheaves, since $S(U)_n^{(red)} \subset S(X)_n^{(red)}$ for any n .

By [8, th. 4.1.1] for any affine scheme U the complex (7) is an exact complex. Therefore we constructed a flasque resolution of the sheaf $\mathcal{F}$ on X . This resolution calculates the cohomology of the sheaf $\mathcal{F}$ on X .

Remark 3 We constructed here reduced adeles, since we used only nondegenerate simplices in $S(X)$. These reduced adeles really carry information and they are the part of the full complex, see [8].

3.3 Adeles on algebraic surfaces.

In this section we verify that the general adelic complex constructed in previous section coincides with the adelic complex for curves from section 3.1. We give also application of general construction of adelic complexes to algebraic surfaces.

We consider a smooth connected algebraic curve C over field k . The set $S(C)_0^{(red)}$ consists of the general point η and all closed points p of the curve C . The set $S(C)_1^{(red)}$ consists of all pairs (η, p) . For any coherent sheaf $\mathcal{F}$ on C we can compute by definition

$$\mathbb{A}_C^0(\mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}_C} k(C) \times \prod_{p \in C} \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p}.$$

Let subset $K \subset S(C)_0^{(red)}$ consists of all closed points of the curve C . We have by definition

$$\begin{aligned} \mathbb{A}_C^1(\mathcal{F}) &= \mathbb{A}(K, \mathcal{F}_\eta) = \mathbb{A}(K, \mathcal{F} \otimes_{\mathcal{O}_C} k(C)) = \\ &= \mathbb{A}(K, \varinjlim_{D \in \text{Div}(C)} \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_C(D)) = \\ &= \varinjlim_{D \in \text{Div}(C)} \mathbb{A}(K, \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_C(D)) = \\ &= \varinjlim_{D \in \text{Div}(C)} \prod_{p \in C} \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p}(D). \end{aligned}$$

Therefore the adelic complex constructed in section 3.2.3 coincides with the adelic complex for curves from section 3.1.

We consider a smooth connected algebraic surface X over field k .

The set $S(X)_0^{(red)}$ consists of the general point η of X , general points of all irreducible curves $C \subset X$, all closed points $p \in X$.

The set $S(X)_1^{(red)}$ consists of all pairs (η, C) , (η, p) and (C, p) . (In our notations we identify the general point of a curve $C \subset X$ with the curve C .)

The set $S(X)_2^{(red)}$ consists of all triples (η, C, p) .

We consider $\delta = (\eta, C, p)$. Let f be a natural map from the local ring $\mathcal{O}_{p,X}$ to the completion $\hat{\mathcal{O}}_{p,X}$. The curve C defines a prime ideal $\mathbf{C}'$ in the ring $\mathcal{O}_{p,X}$. Let $\mathbf{C}_1, \dots, \mathbf{C}_n$ be all prime ideals of height 1 in the ring $\hat{\mathcal{O}}_{p,X}$ such that for any i $f^{-1}(\mathbf{C}_i) = \mathbf{C}'$. Any such $\mathbf{C}_i$ we will call a germ of $\mathbf{C}$ at p . For any such germ $\mathbf{C}_i$ we define a two-dimensional local field

$$K_{p,\mathbf{C}_i} = \text{Frac} \varprojlim_l (\hat{\mathcal{O}}_{p,X(\mathbf{C}_i)} / \mathbf{C}_i^l \hat{\mathcal{O}}_{p,X(\mathbf{C}_i)}).$$

The ring $\hat{\mathcal{O}}_{p,X(\mathbf{C}_i)}$ is a localization of the ring $\hat{\mathcal{O}}_{p,X}$ along the prime ideal $\mathbf{C}_i$.

Then according to formula (3), we have (see [4], [19])

$$\mathbb{A}(\delta, \mathcal{O}_X) = K_\delta = \bigoplus_{i=1}^{i=n} K_{p,\mathbf{C}_i}.$$

Similarly we have

$$\mathbb{A}((C, p), \mathcal{O}_X) = \bigoplus_{i=1}^{i=n} \mathcal{O}_{K_{p,\mathbf{C}_i}}.$$

We compute by definition

$$\mathbb{A}((\eta, C), \mathcal{O}_X) = K_C,$$

where the field K_C is the completion of the field $k(X)$ along the discrete valuation given by irreducible curve C on X .

And from definition we obtain that $\mathbb{A}((\eta, p), \mathcal{O}_X)$ is a subring in $\text{Frac}(\hat{\mathcal{O}}_{p,X})$ generated by subrings $k(X)$ and $\hat{\mathcal{O}}_{p,X}$. We denote this subring by K_p .

By definition we compute

$$\mathbb{A}((\eta), \mathcal{O}_X) = k(X),$$

$$\mathbb{A}((C), \mathcal{O}_X) = \mathcal{O}_{K_C},$$

$$\mathbb{A}((p), \mathcal{O}_X) = \hat{\mathcal{O}}_{p,X}.$$

Remark 4 The local boundary maps d_i^n give natural embeddings of rings $\mathbb{A}((\eta), \mathcal{O}_X)$, $\mathbb{A}((C), \mathcal{O}_X)$, $\mathbb{A}((p), \mathcal{O}_X)$, $\mathbb{A}((\eta, p), \mathcal{O}_X)$, $\mathbb{A}((\eta, C), \mathcal{O}_X)$, $\mathbb{A}((C, p), \mathcal{O}_X)$ to the ring K_δ .

By definition we have

$$\mathbb{A}_X^0(\mathcal{O}_X) = k(X) \times \prod_{C \subset X} \mathcal{O}_{K_C} \times \prod_{p \in X} \hat{\mathcal{O}}_{p,X}.$$

From proposition 1 and similarly to the case of algebraic curve we compute the ring $\mathbb{A}_X^2(\mathcal{O}_X)$, see details in [4], [21].

For any prime ideal $\mathbf{C} \subset \hat{\mathcal{O}}_{p,X}$ of height 1 we define the subring $\hat{\mathcal{O}}_{p,X}(\infty\mathbf{C})$ of $K_{p,\mathbf{C}}$:

$$\hat{\mathcal{O}}_{p,X}(\infty\mathbf{C}) = \varinjlim_l t_{\mathbf{C}}^{-l} \hat{\mathcal{O}}_{p,X},$$

where $t_{\mathbf{C}}$ is a generator of ideal $\mathbf{C}$ in $\hat{\mathcal{O}}_{p,X}$.

The ring $\hat{\mathcal{O}}_{p,X}(\infty\mathbf{C})$ does not depend on the choice of $t_{\mathbf{C}}$.

By $p \in \mathbf{C} \subset X$ we denote a germ at p of an irreducible curve on X . Now we have

$$\mathbb{A}_X^2(\mathcal{O}_X) = \{f_{p,\mathbf{C}}\} \in \prod_{p \in \mathbf{C} \subset X} K_{p,\mathbf{C}} \quad \text{under the following two conditions.}$$

1. There exists a divisor D on X such that for any $p \in \mathbf{C} \subset X$

$$\nu_{K_{p,\mathbf{C}}}(f_{p,\mathbf{C}}) \geq \nu_{\mathbf{C}}(D).$$

2. For any irreducible curve $C \subset X$, any integer k and all except a finite number of points $p \in C$ we have that inside of group $(K_{p,\mathbf{C}} \bmod \mathbf{C}^k \mathcal{O}_{K_{p,\mathbf{C}}})$

$$f_{p,C} \bmod \mathbf{C}^k \mathcal{O}_{K_{p,\mathbf{C}}} \in \hat{\mathcal{O}}_{p,X}(\infty\mathbf{C}) \bmod \mathbf{C}^k \mathcal{O}_{K_{p,\mathbf{C}}}.$$

Here we supposed that the curve C has at p a germ $\mathbf{C}$.

We have

$$\mathbb{A}_X^1(\mathcal{O}_X) = \left(\prod_{C \subset X} K_C \right) \cap \mathbb{A}_X^2(\mathcal{O}_X) \times \left(\prod_{p \in X} K_p \right) \cap \mathbb{A}_X^2(\mathcal{O}_X) \times \left(\prod_{p \in \mathbf{C} \subset X} \mathcal{O}_{K_{p,\mathbf{C}}} \right) \cap \mathbb{A}_X^2(\mathcal{O}_X),$$

where we take the intersection inside of $\prod_{p \in \mathbf{C} \subset X} K_{p,\mathbf{C}}$ due to remark 4 and diagonal embeddings

$$\prod_{C \subset X} K_C \longrightarrow \prod_{p \in \mathbf{C} \subset X} K_{p,\mathbf{C}}$$

and

$$\prod_{p \in X} K_p \longrightarrow \prod_{p \in \mathbf{C} \subset X} K_{p,\mathbf{C}}.$$

From formula 6 and explicit description of rings $\mathbb{A}_X^*(\mathcal{O}_X)$ it is easy to see differentials d^n in the complex $\mathcal{A}_X(\mathcal{O}_X)$ ([4], [21]). Indeed, let

$$A_0 = k(X), \quad A_1 = \prod_{C \subset X} \mathcal{O}_{K_C}, \quad A_2 = \prod_{p \in X} \hat{\mathcal{O}}_{p,X},$$

$$\begin{aligned}
A_{01} &= \left(\prod_{C \subset X} K_C \right) \cap \mathbb{A}_X^2(\mathcal{O}_X), & A_{02} &= \left(\prod_{p \in X} K_p \right) \cap \mathbb{A}_X^2(\mathcal{O}_X), \\
A_{12} &= \left(\prod_{p \in \mathbb{C} \subset X} \mathcal{O}_{K_p, \mathbb{C}} \right) \cap \mathbb{A}_X^2(\mathcal{O}_X), & A_{012} &= \mathbb{A}_X^2(\mathcal{O}_X).
\end{aligned}$$

Then adelic complex $\mathcal{A}_X(\mathcal{O}_X)$ is

$$\begin{array}{ccccc}
A_0 \oplus A_1 \oplus A_2 & \longrightarrow & A_{01} \oplus A_{02} \oplus A_{12} & \longrightarrow & A_{012} \\
(a_0, a_1, a_2) & \mapsto & (a_1 - a_0, a_2 - a_0, a_2 - a_1) & & \\
& & (a_{01}, a_{02}, a_{12}) & \mapsto & a_{01} - a_{02} + a_{12}.
\end{array}$$

Remark 5 We remark the following interesting property, see [24, remark 5], [4]. For any subset $I \subset [0, 1, 2]$ we have an embedding $A_I \hookrightarrow A_{012}$. Now for any subsets $I, J \subset [0, 1, 2]$ we have that inside of group A_{012}

$$A_I \cap A_J = A_{I \cap J}.$$

This property is also true for corresponding components of adelic complex of any locally free sheaf on X .

4 Restricted adelic complexes

In this section we describe restricted adelic complexes. The main difference of restricted adelic complexes from adelic complexes constructed in section 3 is that restricted adelic complexes are connected with one fixed chain (or flag) of irreducible subvarieties of a scheme X .

Restricted adelic complexes come from so-called Krichever correspondence, [24], [17], but see also [23] for connections with the theory of ζ -functions of algebraic curves. Restricted adelic complexes on algebraic curves come originally from the theory of integrable systems, see [25]. For algebraic surfaces restricted adelic complexes were constructed by A.N. Parshin in [24]. For higher dimensional schemes restricted adelic complexes were constructed by author in [17].

4.1 Restricted adelic complexes on algebraic curves and surfaces.

We consider an irreducible algebraic curve C over k . We fix a smooth closed point $p \in C$. For any coherent sheaf $\mathcal{F}$ of rank r on C we consider the following complex

$$\begin{array}{ccc}
\Gamma(C \setminus p, \mathcal{F}) \oplus (\mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p}) & \longrightarrow & \mathcal{F} \otimes_{\mathcal{O}_C} K_p \\
(a_0 \oplus a_1) & \mapsto & a_1 - a_0.
\end{array} \tag{8}$$

We note that for the torsion free sheaf $\mathcal{F}$ we have natural embeddings

$$\mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p} \longrightarrow \mathcal{F} \otimes_{\mathcal{O}_C} K_p$$

$$\Gamma(C \setminus p, \mathcal{F}) \longrightarrow \mathcal{F} \otimes_{\mathcal{O}_C} K_p,$$

where the last embedding is given by

$$\Gamma(C \setminus p, \mathcal{F}) \longrightarrow \Gamma(\mathrm{Spec} \mathcal{O}_p \setminus p, \mathcal{F}) \longrightarrow \Gamma(\mathrm{Spec} \mathcal{O}_{K_p} \setminus p, \mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}_C} K_p.$$

Besides, after the choice of basis of module $\mathcal{F}_p$ over the ring $\mathcal{O}_p$ we have

$$\mathcal{F} \otimes_{\mathcal{O}_C} K_p = K_p^{\oplus r}.$$

Therefore in this case complex (8) is a complex of subgroups inside of $K_p^{\oplus r}$, where K_p is a 1-dimensional local field.

There is the following theorem, see, for example, [24], [23].

Theorem 6 *The cohomology groups of complex (8) coincide with the cohomology groups $H^*(C, \mathcal{F})$.*

Chain of quasi-isomorphisms between complex (8) and adelic complex $\mathcal{A}_C(\mathcal{F})$ was constructed in [24]. It proves the theorem 6. We remark that it is important for the proof, that $C \setminus p$ is an affine curve, see also remark 7 below.

The complex (8) is called *restricted* adelic complex on C associated with the point p .

Now let X be an algebraic surface over k . We fix an irreducible curve $C \subset X$, and a point $p \in C$ which is a smooth point on both C and X . Let $\mathcal{F}$ be a torsion free coherent sheaf on X . We introduce the following notations from [24], [23]. Let $x \in C$,

$$\hat{\mathcal{F}}_x \quad , \quad \hat{\mathcal{F}}_C \quad , \quad \hat{\mathcal{F}}_\eta$$

be completions of stalks of the sheaf $\mathcal{F}$ at scheme points given by x , irreducible curve C and general point η of X correspondingly.

$$B_x(\mathcal{F}) = \bigcap_{\mathbf{D} \neq C} ((\hat{\mathcal{F}}_x \otimes K_x) \cap (\hat{\mathcal{F}}_x \otimes \mathcal{O}_{K_{x,\mathbf{D}}}),$$

where $\mathbf{D}$ runs over all germs at x of irreducible curves on X , which are not equal to C , and the intersection is done inside of the group $\hat{\mathcal{F}}_x \otimes K_x$,

$$B_C(\mathcal{F}) = (\hat{\mathcal{F}}_C \otimes K_C) \cap \left(\bigcap_{x \neq p} B_x \right),$$

where the intersection is done inside of $\hat{\mathcal{F}}_p \otimes K_{x,\mathbf{C}}$ for all closed points $x \neq p$ of C and all germs $\mathbf{C}$ at x of C ,

$$A_C(\mathcal{F}) = B_C(\mathcal{F}) \cap \hat{\mathcal{F}}_C,$$

$$A(\mathcal{F}) = \hat{\mathcal{F}}_\eta \cap \left(\bigcap_{x \in X-C} \hat{\mathcal{F}}_p \right).$$

We note that

$$A(\mathcal{F}) = \Gamma(X - C, \mathcal{F})$$

and for the smooth point $x \in C$ the space $B_x(\mathcal{O}_X)$ coincides with the space $\hat{\mathcal{O}}_{p,X}(\infty\mathbf{C})$ from section 3.3.

The following theorem was proved in [24, th. 3].

Theorem 7 *Let X be an irreducible algebraic surface over a field k , $C \subset X$ be an irreducible curve, and $p \in C$ be a smooth point on both C and X . Let $\mathcal{F}$ be a torsion free coherent sheaf on X . Assume that the surface $X - C$ is affine. Then there exists a chain of quasi-isomorphisms between adelic complex $\mathcal{A}_X(\mathcal{F})$ and the following complex*

$$\begin{array}{ccccc} A(\mathcal{F}) \oplus A_C(\mathcal{F}) \oplus \hat{\mathcal{F}}_p & \longrightarrow & B_C(\mathcal{F}) \oplus B_p(\mathcal{F}) \oplus (\hat{\mathcal{F}}_p \otimes \mathcal{O}_{K_{p,\mathbf{C}}}) & \longrightarrow & \hat{\mathcal{F}}_p \otimes K_{p,\mathbf{C}} \\ (a_0, a_1, a_2) & \mapsto & (a_1 - a_0, a_2 - a_0, a_2 - a_1) & & \\ & & (a_{01}, a_{02}, a_{12}) & \mapsto & a_{01} - a_{02} + a_{12}. \end{array} \quad (9)$$

Under the conditions of this theorem the cohomology groups of the complex (9) coincide with the cohomology groups of adelic complex $\mathcal{A}_X(\mathcal{F})$, and therefore they are equal to $H^*(X, \mathcal{F})$.

Definition 6 *Complex (9) is called restricted adelic complex on X associated with the curve C and the point $p \in C$.*

There is the following proposition, see [24, prop.4].

Proposition 8 *Under the conditions of theorem 7 we suppose also that $\mathcal{F}$ is a locally free sheaf, X is a projective variety, the local rings of X are Cohen-Macaulay and the curve C is locally complete intersection. Then, inside the field $K_{x,\mathbf{C}}$, we have*

$$B_C(\mathcal{F}) \cap B_p(\mathcal{F}) = A(\mathcal{F}).$$

Let the rank of $\mathcal{F}$ be r . Then after the choice of basis of 1-dimensional free $\hat{\mathcal{O}}_{p,X}$ -module $\hat{\mathcal{F}}_p$ we have

$$\begin{aligned} \hat{\mathcal{F}}_p &= \hat{\mathcal{O}}_{p,X}^{\oplus r}, \\ \hat{\mathcal{F}}_p \otimes K_{p,\mathbf{C}} &= K_{p,\mathbf{C}}^{\oplus r}, \\ \hat{\mathcal{F}}_p \otimes \hat{\mathcal{O}}_{K_{p,\mathbf{C}}} &= \hat{\mathcal{O}}_{K_{p,\mathbf{C}}}^{\oplus r}, \\ B_p(\mathcal{F}) &= B_p^{\oplus r} = \hat{\mathcal{O}}_{p,X}(\infty\mathbf{C})^{\oplus r}, \\ A_C(\mathcal{F}) &= A(\mathcal{F}) \cap \hat{\mathcal{O}}_{K_{p,\mathbf{C}}}^{\oplus r}, \end{aligned}$$

where the last intersection is done inside of $K_{p,\mathbf{C}}^{\oplus r}$.

Now due to proposition 8 we obtain that complex (9) is a complex of subgroups of $K_{p,\mathbf{C}}^{\oplus r}$ and is uniquely determined by one subgroup $B_C(\mathcal{F})$ of $K_{p,\mathbf{C}}^{\oplus r}$. In fact, all the other components of complex (9) can be defined by intersections of $B_C(\mathcal{F})$ with subgroups of $K_{p,\mathbf{C}}^{\oplus r}$, which do not really depend on the sheaf $\mathcal{F}$.

4.2 Restricted adelic complexes on higherdimensional schemes.

In this section we construct restricted adelic complexes for arbitrary schemes. These complexes will generalize corresponding complexes from section 4.1.

4.2.1 General definitions.

Let X be a Noetherian separated scheme. Consider a flag of closed subschemes

$$X \supset Y_0 \supset Y_1 \supset \dots \supset Y_n$$

in X . Let J_j be the ideal sheaf of Y_j in X , $0 \leq j \leq n$. Let i_j be the embedding $Y_j \hookrightarrow X$. Let U_i be an open subscheme of Y_i complementing Y_{i+1} , $0 \leq i \leq n-1$. Let $j_i : U_i \hookrightarrow Y_i$ be the open embedding of U_i in Y_i , $0 \leq i \leq n-1$. Put $U_n = Y_n$ and let j_n be the identity morphism from U_n to Y_n .

Assume that every point $x \in X$ has an open affine neighbourhood $U \ni x$ such that $U \cap U_i$ is an affine scheme for any $0 \leq i \leq n$. In what follows, a flag of subschemes $\{Y_i, 0 \leq i \leq n\}$ with this condition is called a flag with *locally affine complements*.

Remark 6 The last condition (existence of locally affine complements) holds, for example, in the following cases:

- Y_{i+1} is the Cartier divisor on Y_i for $0 \leq i \leq n-1$, and
- U_i is an affine scheme for any $0 \leq i \leq n-1$ (the intersection of two open affine subschemes on a separated scheme is an affine subscheme.)

Consider the n -dimensional simplex and its standard simplicial set (without degeneracy). To be precise, consider the set:

$$(\{0\}, \{1\}, \dots, \{n\})$$

(all the integers between 0 and n .) Then the simplicial set $S = \{S_k\}$ is given by

- $S_0 \stackrel{\text{def}}{=} \{\eta \in \{0\}, \{1\}, \dots, \{n\}\}$.
- $S_k \stackrel{\text{def}}{=} \{(\eta_0, \dots, \eta_k), \text{ where } \eta_l \in S_0 \text{ and } \eta_{l-1} < \eta_l\}$.

The boundary map ∂_i ($0 < i < k$) is given by eliminating the i -th component of the vector $(\eta_0, \dots, \eta_k)$ to give the i -th face of $(\eta_0, \dots, \eta_k)$.

Let $\mathbf{QS}(X)$ be the category of quasicoherent sheaves on X . Let $\mathbf{Sh}(X)$ be the category of sheaves of Abelian groups on X . Let $f : Y \longrightarrow X$ be a morphism of schemes. Then f^* always denotes the pull-back functor in the category of sheaves of Abelian groups, and f_* is the direct image functor in the category of sheaves of Abelian groups.

We give the following definition from [17].

Definition 7 For any $(\eta_0, \dots, \eta_k) \in S_k$ we define a functor

$$V_{(\eta_0, \dots, \eta_k)} : \mathbf{QS}(X) \longrightarrow \mathbf{Sh}(X),$$

which is uniquely determined by the following inductive conditions:

1. $V_{(\eta_0, \dots, \eta_k)}$ commutes with direct limits.
2. If $\mathcal{F}$ is a coherent sheaf and $\eta \in S_0$, then

$$V_\eta(\mathcal{F}) \stackrel{\text{def}}{=} \varprojlim_{m \in \mathbf{N}} (i_\eta)_* (j_\eta)^* (\mathcal{F} / J_\eta^m \mathcal{F}).$$

3. If $\mathcal{F}$ is a coherent sheaf and $(\eta_0, \dots, \eta_k) \in S_k$, $k \geq 1$, then

$$V_{(\eta_0, \eta_1, \dots, \eta_k)}(\mathcal{F}) \stackrel{\text{def}}{=} \varprojlim_{m \in \mathbf{N}} V_{(\eta_1, \dots, \eta_k)} \left((i_{\eta_0})_* (j_{\eta_0})^* (\mathcal{F} / J_{\eta_0}^m \mathcal{F}) \right).$$

We'll use sometimes the equivalent notation for $V_{(\eta_0, \dots, \eta_k)}(\mathcal{F})$, in which the closed subschemes are indicated explicitly:

$$V_{(\eta_0, \dots, \eta_k)}(\mathcal{F}) = V_{(Y_{\eta_0}, \dots, Y_{\eta_k})}(X, \mathcal{F}).$$

There is the following proposition, [17, prop. 1], which is proved by induction.

Proposition 9 Let $\sigma = (\eta_0, \dots, \eta_k) \in S_k$. Then the following assertions hold.

1. The functor $V_\sigma : \mathbf{QS}(X) \longrightarrow \mathbf{Sh}(X)$ is well defined.
2. The functor V_σ is exact and additive.
3. The functor V_σ is local on X , that is, for any open $U \subset X$ and any quasicohherent sheaf $\mathcal{F}$ on X we have

$$V_{(Y_{\eta_0}, \dots, Y_{\eta_k})}(X, \mathcal{F})|_U = V_{(Y_{\eta_0} \cap U, \dots, Y_{\eta_k} \cap U)}(U, \mathcal{F}|_U).$$

(if $Y_j \cap U = \emptyset$, then $Y_i \cap U$ is the empty subscheme of U defined by the ideal sheaf $\mathcal{O}_U$.)

4. For any quasicohherent sheaf $\mathcal{F}$ on X the sheaf $V_{(\eta_0, \dots, \eta_k)}(\mathcal{F})$ is a sheaf of $\mathcal{O}_X$ -modules supported on the subscheme Y_{η_k} . (In general, this sheaf is not quasicohherent.)
5. For any quasicohherent sheaf $\mathcal{F}$ on X we have

$$V_\sigma(\mathcal{F}) = V_\sigma(\mathcal{O}_X) \otimes_{\mathcal{O}_X} \mathcal{F}.$$

6. If all U_i is affine, $0 \leq i \leq n$, then for any quasicohherent sheaf $\mathcal{F}$ on X and any $m \geq 1$ we have

$$H^m(X, V_\sigma(\mathcal{F})) = 0.$$

4.2.2 Construction of restricted adelic complex.

We consider the standard n -simplex $S = \{S_k, 0 \leq k \leq n\}$ without degeneracy. If $\sigma = (\eta_0, \dots, \eta_k) \in S_k$, then $\partial_i(\sigma)$ is the i th face of σ , $0 \leq i \leq k$. We define a morphism of functors $d_i(\sigma) : V_{\partial_i(\sigma)} \longrightarrow V_\sigma$, as the morphism that commutes with direct limits and coincides on coherent sheaves with the map

$$V_{\partial_i(\sigma)}(\mathcal{F}) \longrightarrow V_\sigma(\mathcal{F}) \quad (10)$$

defined by the following rules.

- a) If $i = 0$, then (10) is obtained by applying the functor $V_{\partial_0(\sigma)}$ to the map

$$\mathcal{F} \longrightarrow (i_{\eta_0})_*(j_{\eta_0})_*(j_{\eta_0})^*(\mathcal{F}/J_{\eta_0}^m \mathcal{F})$$

and passing to the projective limit with respect to m ;

- b) If $i = 1$ and $k = 1$, then we have the natural map

$$(i_{\eta_0})_*(j_{\eta_0})_*(j_{\eta_0})^*(\mathcal{F}/J_{\eta_0}^m \mathcal{F}) \longrightarrow V_{(\eta_1)}((i_{\eta_0})_*(j_{\eta_0})_*(j_{\eta_0})^*(\mathcal{F}/J_{\eta_0}^m \mathcal{F})).$$

Passing to the projective limit with respect to m , we get the map (10) in this case.

- c) If $i \neq 0$ and $k > 1$, then we use induction on k to get the map

$$V_{\partial_{i-1}(\partial_0(\sigma))}((i_{\eta_0})_*(j_{\eta_0})_*(j_{\eta_0})^*(\mathcal{F}/J_{\eta_0}^m \mathcal{F})) \longrightarrow V_{\partial_0(\sigma)}((i_{\eta_0})_*(j_{\eta_0})_*(j_{\eta_0})^*(\mathcal{F}/J_{\eta_0}^m \mathcal{F})).$$

Passing to the projective limit to m we get the map (10) in this case.

There is the following proposition, [17, prop. 3].

Proposition 10 *For any $1 \leq k \leq n$, $0 \leq i \leq k$ let*

$$d_i^k \stackrel{\text{def}}{=} \sum_{\sigma \in S_k} d_i(\sigma) \quad : \quad \bigoplus_{\sigma \in S_{k-1}} V_\sigma \longrightarrow \bigoplus_{\sigma \in S_k} V_\sigma.$$

We define the map

$$d_0^0 \quad : \quad id \longrightarrow \bigoplus_{\sigma \in S_0} V_\sigma$$

as the direct sum of the natural maps $\mathcal{F} \longrightarrow V_\sigma(\mathcal{F})$. (Here id is the functor of the natural imbedding of $\mathbf{QS}(X)$ into $\mathbf{Sh}(X)$, $\mathcal{F}$ is a quasicoherent sheaf on X , $\sigma \in S_0$.)

Then for all $0 \leq i < j \leq k \leq n-1$ we have

$$d_j^{k+1} d_i^k = d_i^{k+1} d_{j-1}^k. \quad (11)$$

Let

$$d^m \stackrel{\text{def}}{=} \sum_{0 \leq i \leq m} (-1)^i d_i^m$$

Then, given any quasicoherent sheaf $\mathcal{F}$ on X , proposition 10 enables us to construct the complex of sheaves $V(\mathcal{F})$ in the standard way:

$$\dots \longrightarrow \bigoplus_{\sigma \in S_{m-1}} V_{\sigma}(\mathcal{F}) \xrightarrow{d^m} \bigoplus_{\sigma \in S_m} V_{\sigma}(\mathcal{F}) \longrightarrow \dots$$

The property $d^{m+1}d^m = 0$ follows from (11) by an easy direct calculation.

We have the following theorem, [17, th. 1].

Theorem 8 *Let X be a Noetherian separated scheme and let $Y_0 \supset Y_1 \supset \dots \supset Y_n$ be a flag of closed subschemes with locally affine complements. Assume that $Y_0 = X$. Then the following complex is exact:*

$$0 \longrightarrow \mathcal{F} \xrightarrow{d^0} V(\mathcal{F}) \longrightarrow 0. \quad (12)$$

Proof. . We give the sketch of the proof. It suffices to consider the case when the sheaf $\mathcal{F}$ is coherent. We consider the exact sequence of sheaves

$$0 \longrightarrow \mathcal{H} \longrightarrow \mathcal{F} \longrightarrow (j_0)_*(j_0)^*\mathcal{F} \longrightarrow \mathcal{G} \longrightarrow 0.$$

Since the functors V_{σ} are exact for all σ , we obtain the following exact sequence of complexes of sheaves:

$$0 \longrightarrow V(\mathcal{H}) \longrightarrow V(\mathcal{F}) \longrightarrow V((j_0)_*(j_0)^*\mathcal{F}) \longrightarrow V(\mathcal{G}) \longrightarrow 0. \quad (13)$$

The sheaves $\mathcal{H}$ and $\mathcal{G}$ are supported on Y_1 . Therefore by induction we may assume that the complexes

$$\begin{aligned} 0 \longrightarrow \mathcal{H} &\xrightarrow{d^0} V(\mathcal{H}) \longrightarrow 0, \\ 0 \longrightarrow \mathcal{G} &\xrightarrow{d^0} V(\mathcal{G}) \longrightarrow 0 \end{aligned}$$

are already exact. The complex

$$0 \longrightarrow (j_0)_*(j_0)^*\mathcal{F} \xrightarrow{d^0} V((j_0)_*(j_0)^*\mathcal{F}) \longrightarrow 0$$

is exact, because the complex $V((j_0)_*(j_0)^*\mathcal{F})$ has the same components $V_{\sigma'}(\mathcal{F})$ of degrees k and $k+1$, $\sigma' = (0, \eta_0, \dots, \eta_k)$ for $\sigma = (\eta_0, \dots, \eta_k) \in S_k$. Now the theorem follows from sequence (13).

For any $\sigma \in S_k$ we define

$$A_{\sigma}(\mathcal{F}) \stackrel{\text{def}}{=} H^0(X, V_{\sigma}(\mathcal{F})).$$

We have the following proposition, [17, prop. 4].

Proposition 11 *Let X be a Noetherian separated scheme, and let $Y_0 \supset Y_1 \supset \dots \supset Y_n$ be a flag of closed subschemes such that U_i is affine, $0 \leq i \leq n$. Let $\sigma \in S_k$ be arbitrary.*

1. Then A_σ is an exact and additive functor: $\mathbf{QS}(X) \longrightarrow \mathbf{Ab}$.
2. If $X = \text{Spec } A$ and M is some A -module, then

$$A_\sigma(\tilde{M}) = A_\sigma(\mathcal{O}_X) \otimes_A M.$$

Let $\mathcal{F}$ be any quasicoherent sheaf on X . Applying the functor $H^0(X, \cdot)$ to the complex $V(\mathcal{F})$, we obtain the complex $A(\mathcal{F})$ of Abelian groups:

$$\dots \longrightarrow \bigoplus_{\sigma \in S_{m-1}} A_\sigma(\mathcal{F}) \longrightarrow \bigoplus_{\sigma \in S_m} A_\sigma(\mathcal{F}) \longrightarrow \dots$$

Now we have the following theorem, see [17, th. 2].

Theorem 9 *Let X be a Noetherian separated scheme. Let $Y_0 \supset Y_1 \supset \dots \supset Y_n$ be a flag of closed subschemes such that $Y_0 = X$ and U_i is affine, $0 \leq i \leq n$. Then the cohomology of the complex $A(\mathcal{F})$ coincide with that of the sheaf $\mathcal{F}$ on X , that is, for any i*

$$H^i(X, \mathcal{F}) = H^i(A(\mathcal{F})).$$

Proof. It follows from theorem 8 and assertion 6 of proposition 9 that $V(\mathcal{F})$ is an acyclic resolution for the sheaf $\mathcal{F}$. Hence the cohomology of $\mathcal{F}$ may be calculated by means of global sections of this resolution. Theorem 9 is proved.

This theorem immediatly yields the following geometric corollary, see [17, th. 3].

Theorem 10 *Let X be a projective algebraic scheme of dimension n over a field. Let $Y_0 \supset Y_1 \supset \dots \supset Y_n$ be a flag of closed subschemes such that $Y_0 = X$ and Y_i is an ample divisor on Y_{i-1} for $1 \leq i \leq n$. Then for any quasicoherent sheaf $\mathcal{F}$ on X and any i we have*

$$H^i(X, \mathcal{F}) = H^i(A(\mathcal{F})).$$

Proof. Since Y_i is an ample divisor on Y_{i-1} for $1 \leq i \leq n$, we see that U_i is an affine scheme for all $0 \leq i \leq n-1$. Since $\dim Y_n = 0$, we obtain that $U_n = Y_n$ is also affine. Applying theorem 9 we complete the proof.

Remark 7 For any quasicoherent sheaf $\mathcal{F}$ and any $\sigma = (\eta_0) \in S_0$, $A_\sigma(\mathcal{F})$ is the group of section over U_{η_0} of the sheaf $\mathcal{F}$ lifted to the formal neighbourhood of the subscheme Y_{η_0} in X . The complex $A(\mathcal{F})$ can be interpreted as the Čech complex for this "acyclic covering" of the scheme X .

Definition 8 *The complex $A(\mathcal{F})$ is called restricted adelic complex on X associated with flag $Y_0 \supset Y_1 \supset \dots \supset Y_n$.*

Remark 8 • If C is an algebraic curve, $Y_0 = C$ and $Y_1 = p$ is a smooth point, then $A(\mathcal{F})$ coincides with complex (8). Indeed,

$$A_0(\mathcal{F}) = \Gamma(C \setminus p, \mathcal{F}),$$

$$A_1(\mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}_C} \mathcal{O}_{K_p},$$

$$A_{12}(\mathcal{F}) = \mathcal{F} \otimes_{\mathcal{O}_C} K_p.$$

- If X is an algebraic surface, $Y_0 = C$ and $Y_1 = p$ is a smooth point on both C and X , then $A(\mathcal{F})$ coincides with complex (9). Indeed,

$$A_0(\mathcal{F}) = A(\mathcal{F}), \quad A_1(\mathcal{F}) = A_C(\mathcal{F}),$$

$$A_2(\mathcal{F}) = \hat{\mathcal{F}}_p, \quad A_{01}(\mathcal{F}) = B_C(\mathcal{F}),$$

$$A_{02}(\mathcal{F}) = B_p(\mathcal{F}), \quad A_{12}(\mathcal{F}) = \hat{\mathcal{F}}_p \otimes \mathcal{O}_{K_{p,\mathbf{C}}},$$

$$A_{012}(\mathcal{F}) = \hat{\mathcal{F}}_p \otimes K_{p,\mathbf{C}}.$$

4.2.3 Reconstruction of restricted adelic complex.

By proposition 10 we have the natural map

$$d_i(\sigma) : A_{\partial_i(\sigma)}(\mathcal{F}) \longrightarrow A_\sigma(\mathcal{F})$$

for any $\sigma \in S_k$, $1 \leq k \leq n$, and any i , $0 \leq i \leq k$.

Taking (11) into account, we obtain the natural map

$$A_{\sigma_1}(\mathcal{F}) \longrightarrow A_{\sigma_2}(\mathcal{F})$$

for any locally free sheaf $\mathcal{F}$ on X and any $\sigma_1, \sigma_2 \in S$, $\sigma_1 \subset \sigma_2$.

There is the following proposition, see [17, th. 4].

Proposition 12 *Let X be a projective equidimensional Cohen-Macaulay scheme of dimension n over a field. Let $Y_0 \supset Y_1 \supset \dots \supset Y_n$ be a flag of closed subschemes such that $Y_0 = X$ and Y_i is an ample Cartier divisor on Y_{i-1} for $1 \leq i \leq n$. Then the following assertions hold for any locally free sheaf $\mathcal{F}$ on X .*

1. *The natural map $H^0(X, \mathcal{F}) \longrightarrow A_\sigma(\mathcal{F})$ is an embedding for any $\sigma \in S_k$, $0 \leq k \leq n$.*
2. *The natural map $A_{\sigma_1}(\mathcal{F}) \longrightarrow A_{\sigma_2}(\mathcal{F})$ is an embedding for any locally free sheaf $\mathcal{F}$ on X and any $\sigma_1, \sigma_2 \in S$, $\sigma_1 \subset \sigma_2$.*

Remark 9 We note that any integral Noetherian scheme of dimension 1 is a Cohen-Macaulay scheme. Any normal Noetherian scheme of dimension 2 is a Cohen-Macaulay scheme, see [7, Ch. II, th. 8.22A].

By $(0, 1, \dots, n) \in S$ we denote the unique face of dimension n in S .

By proposition 12 we can embed $A_{\sigma_1}(\mathcal{F})$ and $A_{\sigma_2}(\mathcal{F})$ in $A_{(0,1,\dots,n)}(\mathcal{F})$ for any $\sigma_1, \sigma_2 \in S$ and any locally free sheaf $\mathcal{F}$.

Now we formulate the following theorem, see [17, th. 5].

Theorem 11 *Let all the hypotheses of proposition 12 be satisfied. Then the following assertions hold for any locally free sheaf $\mathcal{F}$ and any $\sigma_1, \sigma_2 \in S$.*

1. *If $\sigma_1 \cap \sigma_2 = \emptyset$, then, inside of $A_{(0,1,\dots,n)}$,*

$$A_{\sigma_1}(\mathcal{F}) \cap A_{\sigma_2}(\mathcal{F}) = H^0(X, \mathcal{F}).$$

2. *If $\sigma_1 \cap \sigma_2 \neq \emptyset$, then, inside of $A_{(0,1,\dots,n)}$,*

$$A_{\sigma_1}(\mathcal{F}) \cap A_{\sigma_2}(\mathcal{F}) = A_{\sigma_1 \cap \sigma_2}(\mathcal{F}).$$

Remark 10 Theorem 11 is similar to the property of adelic complexes $\mathcal{A}_X(\mathcal{F})$ noticed in remark 5.

We assume that the hypotheses of proposition 12 hold and that the field k of definition of the scheme X is k . We also assume that $Y_n = p$, where p is a smooth point on each Y_i , $0 \leq i \leq n$.

Let us choose and fix local parameters $t_1, \dots, t_n \in \hat{\mathcal{O}}_{p,X}$ such that $t_i|_{Y_{i-1}} = 0$ is a local equation of the divisor Y_i in the formal neighbourhood of the point p on the scheme Y_{i-1} , $1 \leq i \leq n$.

Let $\mathcal{F}$ be a rank 1 locally free sheaf on X . We fix a trivialization e_p of $\mathcal{F}$ in a formal neighbourhood of the point p on X , that is an isomorphism

$$e_p : \hat{\mathcal{F}}_p \longrightarrow \hat{\mathcal{O}}_{p,X}.$$

By the choice of the local parameters and the trivialization we can identify $A_{(0,1,\dots,n)}(\mathcal{F})$ with the n -dimensional local field $k(p)((t_n)) \dots ((t_1))$:

$$A_{(0,1,\dots,n)}(\mathcal{F}) = k(p)((t_n)) \dots ((t_1)).$$

Moreover, we fix a set of integers $0 \leq j_1 \leq \dots \leq j_k \leq n-1$. Define $\sigma_{(j_1,\dots,j_k)} \in S_{n-k}$ as the following set

$$\{i : 0 \leq i \leq n, i \neq j_1, \dots, i \neq j_k\}.$$

By proposition 12 we have a natural embedding

$$A_{\sigma_{(j_1,\dots,j_k)}}(\mathcal{F}) \longrightarrow A_{(0,1,\dots,n)}(\mathcal{F}).$$

When we identify $A_{(0,1,\dots,n)}(\mathcal{F})$ with the field $k(p)((t_n)) \dots ((t_1))$, the space $A_{\sigma_{(j_1,\dots,j_k)}}(\mathcal{F})$ corresponds to the following k -subspace in $k(p)((t_n)) \dots ((t_1))$:

$$\left\{ \sum a_{i_1,\dots,i_n} t_n^{i_n} \dots t_1^{i_1} : a_{i_1,\dots,i_n} \in k(p), i_{j_1+1} \geq 0, i_{j_2+1} \geq 0, \dots, i_{j_k+1} \geq 0 \right\}. \quad (14)$$

Thus, by theorem 11, to determine the images of $A_\sigma(\mathcal{F})$ in $k(p)((t_n)) \dots ((t_1))$ (for any $\sigma \in S$), it suffices to know only one image of $A_{(0,1,\dots,n-1)}$ in $k(p)((t_n)) \dots ((t_1))$. (All the others are obtained by taking the intersection of the image of this one with the standard subspaces (14) of $k(p)((t_n)) \dots ((t_1))$.)

It is clear that these arguments are generalized immediately to locally free sheaves $\mathcal{F}$ of rank r and to the spaces $k(p)((t_n)) \dots ((t_1))^{\oplus r}$.

These arguments lead to the following theorem, which enables to reconstruct restricted adelic complex $A(\mathcal{F})$, see also [17, th. 6].

Theorem 12 *Let all the hypotheses of proposition 12 be satisfied. We also assume that $Y_n = p$, where p is a smooth point on each Y_i , $0 \leq i \leq n$. Let $\mathcal{F}$ be a locally free sheaf on X . Then the subspace*

$$A_{(0,1,\dots,n-1)}(\mathcal{F}) \subset A_{(0,1,\dots,n)}(\mathcal{F})$$

uniquely determines the restricted adelic complex $A(\mathcal{F})$.

5 Reciprocity laws.

Let L be a field of discrete valuation ν_L with the valuation ring $\mathcal{O}_L$ and the maximal idela m_L . Then there is the tame symbol

$$(f, g)_L = (-1)^{\nu_L(f)\nu_L(g)} \frac{f^{\nu_L(g)}}{g^{\nu_L(f)}} \bmod m_L, \quad (15)$$

where f, g are from L^* .

There is the following reciprocity law, see, for example, [26].

Proposition 13 *Let C be a complete smooth algebraic curve over k . For any $f, g \in k(C)^*$ in the following product only a finite number of terms is not equal to 1 and*

$$\prod_{p \in C} \text{Nm}_{k(p)/k} (f, g)_{K_p} = 1, \quad (16)$$

where the product is taken over all closed points $p \in C$, and Nm is the norm map.

Let K be a 2-dimensional local field with the last residue field k . We have a discrete valuation of rank 2 on K :

$$(\nu_1, \nu_2) : K^* \rightarrow \mathbb{Z} \oplus \mathbb{Z}.$$

Here $\nu_1 = \nu_K$ is the discrete valuation of the field K , and

$$\nu_2(b) \stackrel{\text{def}}{=} \nu_{\bar{K}}(\overline{bt_1^{-\nu_1(b)}}),$$

where $\nu_1(t_1) = 1$. We note that ν_2 depends on the choice of local parameter t_1 .

Let m_K be the maximal ideal of $\mathcal{O}_K$, and $m_{\bar{K}}$ be the maximal ideal of $\mathcal{O}_{\bar{K}}$.

We define a map:

$$\nu_K(,) : K^* \times K^* \longrightarrow \mathbb{Z}$$

as the composition of maps:

$$K^* \times K^* \longrightarrow K_2(K) \xrightarrow{\partial_2} \bar{K}^* \xrightarrow{\partial_1} \mathbb{Z},$$

where ∂_i is the boundary map in algebraic K -theory. The map ∂_2 coincides with tame symbol (15) with respect to discrete valuation ν_1 . The map ∂_1 coincides with the discrete valuation $\nu_{\bar{K}}$.

We define a map:

$$(\ , \ , \)_K : K^* \times K^* \times K^* \longrightarrow k^*$$

as the composition of maps

$$K^* \times K^* \times K^* \longrightarrow K_3^M(K) \xrightarrow{\partial_3} K_2(\bar{K}) \xrightarrow{\partial_2} k^*,$$

where K_3^M is the Milnor K -group.

There are the following explicit expressions for these maps (see [4]):

$$\nu_K(f, g) = \nu_1(f)\nu_2(g) - \nu_2(f)\nu_1(g)$$

$$(f, g, h)_K = \text{sign}_K(f, g, h) f^{\nu_K(g, h)} g^{\nu_K(h, f)} h^{\nu_K(f, g)} \bmod m_K \bmod m_{\bar{K}}$$

$$\text{sign}_K(f, g, h) = (-1)^B,$$

where $B = \nu_1(f)\nu_2(g)\nu_2(h) + \nu_1(g)\nu_2(f)\nu_2(h) + \nu_1(h)\nu_2(g)\nu_2(f) + \nu_2(f)\nu_1(g)\nu_1(h) + \nu_2(g)\nu_1(f)\nu_1(h) + \nu_2(h)\nu_1(f)\nu_1(g)$.

Proposition 14 *For any $f, g, h \in K^*$*

$$\text{sign}_K(f, g, h) = (-1)^A, \text{ where}$$

$$A = \nu_K(f, g)\nu_K(f, h) + \nu_K(f, g)\nu_K(f, h) + \nu_K(f, g)\nu_K(f, h) + \nu_K(f, g)\nu_K(f, h)\nu_K(g, h).$$

Proof. It follows from direct calculations modulo 2 with A and B using the explicit expressions above.

Let X be a smooth algebraic surface over k . We recall the following notations from section 3.3. If C is a curve on X , then K_C is the completion of the field $k(X)$ along the discrete valuation given by the irreducible curve C on X . If p is a point on X , then K_p is a subring in $\text{Frac}(\hat{\mathcal{O}}_{p, X})$ generated by subrings $k(X)$ and $\hat{\mathcal{O}}_{p, X}$.

There are the following reciprocity laws, see [4].

Theorem 13 *1. We fix a point $p \in X$ and any $f, g, h \in K_p^*$. Then in the following sum only a finite number of terms is not equal to 0 and*

$$\sum_{X \supset C \ni p} \nu_{K_{p, C}}(f, g) = 0.$$

In the following product only a finite number of terms is not equal to 1 and

$$\prod_{X \supset C \ni p} (f, g, h)_{K_{p, C}} = 1,$$

where the sum and the product are taken over all germs of irreducible curves on X at p .

2. We fix an irreducible projective curve C on X and any $f, g, h \in K_C^*$. Then in the following sum only a finite number of terms is not equal to 0 and

$$\sum_{p \in \mathbf{C} \subset X} [k(p) : k] \cdot \nu_{K_{p, \mathbf{C}}}(f, g) = 0.$$

In the following product only a finite number of terms is not equal to 1 and

$$\prod_{p \in \mathbf{C} \subset X} \text{Nm}_{k(p)/k}(f, g, h)_{K_{p, \mathbf{C}}} = 1,$$

where the sum and the product are taken over all points $p \in C$ and all germs of irreducible curve C at p .

Remark 11 The relative reciprocity laws were constructed in [15] (see also [16] for a short exposition) for a smooth projective morphism f of a smooth algebraic surface X to a smooth algebraic curve S when $\text{char } k = 0$. If $p \in \mathbf{C} \subset X$ then explicit formulas were constructed in [15] for maps

$$K_2(K_{p, \mathbf{C}}) \longrightarrow K_{f(p)}^*.$$

Remark 12 The map $\nu_K(,)$ for 2-dimensional local field K was interpreted in [18] as the commutator of liftings of elements $f, g \in K^*$ in a central extension of group K^* by $\mathbb{Z}$. From this interpretation the reciprocity laws for $\nu_K(,)$ were proved. The proof in [18] uses adelic rings on an algebraic surface X . This is an abstract version of reciprocity law for $\nu_K(,)$ like abstract version of reciprocity law (16) for a projective curve in [1] (and in [28] for the residues of differentials on a projective curve).

Remark 13 We don't describe the reciprocity laws for residues of differentials of 2-dimensional local fields, which were formulated and proved in [19], see also [29].

Remark 14 The symbols $\nu_K(,)$ and $(, ,)_K$ correspond to the non-ramified and tame ramified extensions of 2-dimensional local fields when the last residue field is finite, see [22].

References

- [1] Arbarello E., De Concini C., Kac V. G. *The Infinite Wedge Representation and the Reciprocity Law for Algebraic Curves*, Comm. Math. Phys., **117** (1988), 1-36.
- [2] Beilinson A.A. Residues and Adeles, *Funct. Anal. Pril.*, **14** (1980), no. 1, 44-45; English transl. in *Func. Anal. Appl.*, **14**, no. 1 (1980), 34-35.
- [3] Fimmel T. *Verdier duality for systems of coefficients over simplicial sets*. Math. Nachr. 190 (1998), 51-122
- [4] Fimmel T., Parshin A.N. *Introduction to higher adelic theory*, preprint 1999.

- [5] Gorchinskiy S.O., Parshin A.N. *Adelic Lefschetz formula for the action of a one-dimensional torus*, Trudy Sankt-Peterburgskogo mat. obtshestva, april 2005, 11, p. 37-57 (russian); see also math.AG/0408058.
- [6] Grothendieck A., Dieudonne J. *Elements de geometrie algebrique I*, Springer-Verlag, Berlin, 1971.
- [7] Hartshorne R. *Algebraic Geometry*, GTM 52, Springer 1977.
- [8] Huber A. On the Parshin-Beilinson Adeles for Schemes, Abh. Math. Sem. Univ. Hamburg, 61 (1991), 249-273.
- [9] Huber A. Adele für Schemata und Zariski-Kogomologie, Schriftenreihe des Math. Inst. der Univ. Muenster, 3. Serie, Heft3, 1991.
- [10] Hübl R., Yekutieli A. *Adelic Chern forms and applications*, Amer. J. Math. 121 (1999), no. 4, 797-839.
- [11] Kato K., Saito S. *Global class field theory of arithmetical schemes*, Cont. Math. Vol. 55, Part I (1986), 255-332.
- [12] Lomadze V. G. *On the intersection index of divisors*, Izv. Akad. Nauk SSSR Ser. Mat. **44** (1980), 1120-1130; English transl. in Math. USSR Izv. **17** (1981).
- [13] Lomadze V.G. *On residues in algebraic geometry*, Izv. Akad. Nauk SSSR Ser. Mat. 45(6) (1981), 1258-1287; English transl. in Math. USSR Izv. 19 (1982).
- [14] Matsumura H. *Commutative Algebra*, Benjamin/Cummings, Reading, Mass., 1980.
- [15] Osipov, D. V. *Adelic constructions of direct images of differentials and symbols*. (Russian) Mat. Sb. 188 (1997), no. 5, 59-84; translation in Sb. Math. 188 (1997), no. 5, 697-723, see also math.AG/9802112.
- [16] Osipov D. *Adelic constructions for direct images of differentials and symbols*. Invitation to higher local fields (Münster, 1999), 215-221 (electronic), Geom. Topol. Monogr., 3, Geom. Topol. Publ., Coventry, 2000, see also math.NT/0012152.
- [17] Osipov D. V. *The Krichever correspondence for algebraic varieties*, Izv. Ross. Akad. Nauk Ser. Mat. 65 (2001), no. 5, 91-128; english translation in Izv. Math. 65 (2001), no. 5, 941-975, see also math.AG/0003188.
- [18] Osipov D.V. *Central extensions and reciprocity laws on algebraic surfaces*, 26 pp., to appear in Sb. Math., see also math.NT/0501155.
- [19] Parshin A. N. *On the arithmetic of two-dimensional schemes, I. Repartitions and residues*, Izv. Akad. Nauk SSSR Ser. Mat. **40** (1976), no. 4, 736-773; English transl. in Math. USSR Izv. **10** (1976), no. 4, 695-729.

- [20] Parshin A.N. *Abelian coverings of arithmetical schemes*, DAN USSR, v.243 (1978), 855-858; English transl. in Soviet. Math. Doklady 19 (1978).
- [21] Parshin A. N. *Chern classes, adeles and L-functions*, J. reine angew Math. **341** (1983), 174-192.
- [22] Parshin A. N. *Local class field theory*, Trud. Mat. Inst. AN SSSR, 165 (1984), 143-170; English transl. in Proceedings of the Steklov Institute of Mathematics 1985, Issue 3.
- [23] Parshin A. N. *Higher-dimensional local fields and L-functions*, Invitation to higher local fields (Münster, 1999), 199–213 (electronic), Geom. Topol. Monogr., 3, Geom. Topol. Publ., Coventry, 2000, see also math.NT/0012151.
- [24] Parshin A. N. *Integrable systems and local fields*, Communications in Algebra, 29(9), 4157-4181 (2001), the short version see in math.AG/9911097.
- [25] Segal G., Wilson G. Loop groups and equations of the KdV type, Publ. Math. IHES, 80(1985), 301-342.
- [26] Serre J.-P. *Groupes algebriques et corps de classes*, Hermann, 1959.
- [27] Serre J.-P. *Local fields*, Graduate Texts in Mathematics, 67. Springer-Verlag, New York-Berlin, 1979.
- [28] Tate J. *Residues of differentials on curves*, Ann. scient. Ec. Norm. Sup., 4 serie, t.1, 1968, p.149.
- [29] Yekutieli A. *An explicit construction of the Grothendieck residue complex*, Asterisque 208 (1992).
- [30] Zhukov I.B. *Structure theorems for complete fields*, Proceedings of the St. Petersburg Mathematical Society, Vol. III, 175–192, Amer. Math. Soc. Transl. Ser. 2, 166, Amer. Math. Soc., Providence, RI, 1995.
- [31] Zhukov I.B. *Higher dimensional local fields*. Invitation to higher local fields (Münster, 1999), 5–18 (electronic), Geom. Topol. Monogr., 3, Geom. Topol. Publ., Coventry, 2000, see also math.NT/0012132.

Steklov Mathematical Institute,
 Gubkina str. 8,
 119991, Moscow, Russia
 e-mail d_osipov@mi.ras.ru